

Boundary Manifolds of Line Arrangements

Eriko Hironaka *

February 5, 2008

Abstract

In this paper we describe the complement of real line arrangements in the complex plane in terms of the boundary three-manifold of the line arrangement. We show that the boundary manifold of any line arrangement is a graph manifold with Seifert fibered vertex manifolds, and depends only on the incidence graph of the arrangement. When the line arrangement is defined over the real numbers, we show that the homotopy type of the complement is determined by the incidence graph together with orderings on the edges emanating from each vertex.

1 Introduction

Let $\mathcal{L}$ be a finite union of lines in the complex plane $\mathbb{C}^2$. Two line arrangements $\mathcal{L}_1$ and $\mathcal{L}_2$ are said to be *topologically equivalent* if there is a homeomorphism of pairs

$$(\mathbb{C}^2, \mathcal{L}_1) \rightarrow (\mathbb{C}^2, \mathcal{L}_2).$$

The *incidence graph* $\Gamma_{\mathcal{L}}$ associated to $\mathcal{L}$ is the bipartite graph with line-vertices corresponding to the set of lines $\mathcal{A}$ in $\mathcal{L}$, point-vertices corresponding to the set of points of intersection $\mathcal{P}$ on $\mathcal{L}$ and edges $e(p, L)$ and $e(L, p)$ whenever $p \in L$. A morphism between incidence graphs is a morphism of graphs preserving the vertex labelings, so that line-vertices go to line-vertices and point-vertices go to point-vertices. Two line arrangements $\mathcal{L}_1$ and $\mathcal{L}_2$ are said to be *combinatorially equivalent* if there is an isomorphism of incidence matrices $\Gamma_{\mathcal{L}_1} \rightarrow \Gamma_{\mathcal{L}_2}$.

Our motivation is to understand, given a line arrangement $\mathcal{L}$ in the complex plane $\mathbb{C}^2$, to what extent the topology of the pair $(\mathbb{C}^2, \mathcal{L})$ is determined by the combinatorics of $\mathcal{L}$. It is known that the cohomology of the complement $E_{\mathcal{L}} = \mathbb{C}^2 \setminus \mathcal{L}$ only depends on the incidence graph $\Gamma_{\mathcal{L}}$ [O-T], [G-M], while the homotopy type and fundamental group of $E_{\mathcal{L}}$ depend on more information [Ryb].

In this paper, we describe the homotopy type of the complement $E_{\mathcal{L}}$, when $\mathcal{L}$ is defined over the real numbers, in terms of the boundary 3-manifold $M_{\mathcal{L}}$ of a regular neighborhood

*Research partially supported by N.S.E.R.C. grant OGP0170260

of $\mathcal{L}$ in $\mathbb{C}^2$. We do this by describing $M_{\mathcal{L}}$ and $E_{\mathcal{L}}$ in terms of a graph of manifolds over the incidence graph $\Gamma_{\mathcal{L}}$.

Let $\Gamma = \{\mathcal{V}, \mathcal{E}\}$ be a directed graph, with vertices $\mathcal{V}$ and edges $\mathcal{E}$. Assume that for each oriented edge $e \in \mathcal{E}$ its opposite $\bar{e}$ is also contained in $\mathcal{E}$. Denote by $i(e)$ the initial point of e and $t(e)$ the terminal point of e . Thus, $i(e) = t(\bar{e})$ and $t(e) = i(\bar{e})$. A *graph manifold* $\{M_v, M_e, \phi_e\}$ over a directed graph Γ is a collection of connected vertex manifolds M_v , for $v \in \mathcal{V}$, connected edge manifolds M_e , with $M_e = M_{\bar{e}}$, for $e \in \mathcal{E}$, and inclusions $\phi_e : M_e \rightarrow M_{t(e)}$ which are isomorphisms onto a boundary component of $M_{t(e)}$, and which induce endomorphisms on fundamental groups (cf. [Wal]). A *morphism* of graph manifolds over a given graph Γ is a collection of continuous maps between corresponding vertex and edge manifolds which commute with the maps ϕ_e . The manifold associated to a graph manifold is the space obtained by gluing the M_v together along their boundary components according to the maps ϕ_e .

Theorem 1 *If $\mathcal{L}$ is a complex line arrangement, and $M_{\mathcal{L}}$ is the boundary 3-manifold of $\mathcal{L}$, then $M_{\mathcal{L}}$ has the structure of a graph manifold over $\Gamma_{\mathcal{L}}$ whose vertex manifolds are Seifert fibered 3-manifolds. Furthermore, if $\mathcal{L}_1$ and $\mathcal{L}_2$ are two line arrangements and*

$$\alpha : \Gamma_{\mathcal{L}_1} \rightarrow \Gamma_{\mathcal{L}_2},$$

is an isomorphism of incidence graphs, then there is a compatible isomorphism

$$\beta : M_{\mathcal{L}_1} \rightarrow M_{\mathcal{L}_2}$$

of graph manifolds.

Any decomposition of a 3-manifold by incompressible surfaces gives rise to a presentation of the manifold as a graph manifold. Let H_d be a thickened d -component oriented Hopf link in S^3 , where each pair of component links having linking number 1. Then $S^3 \setminus H_d$ is a 3-manifold with boundary 2-tori and each boundary torus has a natural framing (μ, λ) . Theorem 1 is a consequence of the following more explicit description of $M_{\mathcal{L}}$. The omitted case is treated separately in the beginning of Section 2.

Theorem 2 *Let $\mathcal{L}$ be any line arrangement in $\mathbb{C}^2$ so that each line $L \in \mathcal{A}$ contains at least one point in $\mathcal{P}$. Then the boundary 3-manifold $M_{\mathcal{L}}$ has a torus decomposition into pieces*

$$M_v = S^3 \setminus H_d,$$

where v ranges over vertices in $\Gamma_{\mathcal{L}}$. If v is a point-vertex, there is a one-to-one correspondence between the d boundary components of M_v and the edges emanating from v in $\Gamma_{\mathcal{L}}$. If v is a line-vertex, there is a one-to-one correspondence between $d - 1$ of the d boundary components of M_v and the edges emanating from v in $\Gamma_{\mathcal{L}}$. Given an edge $e = e(p, L)$, the corresponding boundary components $T_{L,p} \subset M_{v_p}$ and $T_{p,L} \subset M_{v_L}$ are identified by ϕ_e and $\phi_{\bar{e}}$ according to the relation:

$$\begin{aligned} \mu_{p,L} &= \lambda_{L,p} \\ \lambda_{p,L} &= \mu_{L,p} + \lambda_{L,p}, \end{aligned}$$

where $(\mu_{L,p}, \lambda_{L,p})$ is the induced framing of $T_{L,p}$ and $(\mu_{p,L}, \lambda_{p,L})$ is the induced framing of $T_{p,L}$.

Theorem 2, proved in Section 2, implies that the fundamental group of $M_{\mathcal{L}}$ is torsion free and residually finite [Ser], [Hemp] (see Corollary 2.4).

Given two topological space $X \subset Y$, let Y/X be the space Y with X collapsed to a point. For real line arrangements, the homotopy type of the complement can be described as follows.

Theorem 3 *Let $\mathcal{L}$ be a real line arrangement. Then there is a continuous map*

$$f : \Gamma_{\mathcal{L}} \rightarrow M_{\mathcal{L}}$$

such that $f(v) \in M_v$ for all vertices v of $\Gamma_{\mathcal{L}}$, and the complement $E_{\mathcal{L}} = \mathbb{C}^2 \setminus \mathcal{L}$ is homotopy equivalent to $M_{\mathcal{L}}/f(\Gamma_{\mathcal{L}})$.

The map f of Theorem 3 depends on the *ordered incidence graph* $\widetilde{\Gamma}_{\mathcal{L}}$ of $\mathcal{L}$, the incidence graph $\Gamma_{\mathcal{L}}$ together with an ordering of the edges emanating from each vertex. (An ordered graph is similar to a *fat graph*, which has a cyclic ordering on the edges at each vertex [Pen].) A morphism of ordered incidence graphs is a morphism of incidence graphs which preserves orderings of edges. Let $\mathcal{L}$ be a line arrangement defined over the real numbers. We construct a model for the homotopy type of $E_{\mathcal{L}}$ which only depends on the ordered graph $\widetilde{\Gamma}_{\mathcal{L}}$ associated to $\mathcal{L}$ and is simple to describe (cf. [Falk], [Lib], [O-T], [Ran] and [Sal].)

Theorem 4 *Let $\mathcal{L}$ be a real line arrangement. Choose basepoints $b_v \in M_v$ for each vertex v of $\Gamma_{\mathcal{L}}$. For each edge e in $\Gamma_{\mathcal{L}}$, let τ_e be a path on $M_{i(e)} \cup M_{t(e)}$ from $b_{i(e)}$ to $b_{t(e)}$ which intersects M_e in one point. Identify $\Gamma_{\mathcal{L}}$ with its associated singular 1-complex. Then there is a continuous mapping*

$$f : \Gamma_{\mathcal{L}} \rightarrow M_{\mathcal{L}}$$

satisfying the following conditions:

- (i) $f(v) = b_v$ for all vertices $v \in \mathcal{V}$;
- (ii) if $p_1, \dots, p_s$ are the ordered points on $L \cap \mathcal{P}$, $e = e(L, p)$, and $p = p_j$, then

$$f(e) = \begin{cases} \tau_e & \text{if } j = 1, 2 \\ g_{L,p_2} \cdots g_{L,p_{j-1}} \tau_e & \text{if } j > 2, \end{cases}$$

where $L_1, \dots, L_r$ are the ordered lines through p , and

$$g_{L_j,p} = \mu_{L_1,p} \cdots \mu_{L_j,p} \mu_{L_{j-1},p}^{-1} \cdots \mu_{L_1,p}^{-1};$$

- (iii) $f(e) = f(\bar{e})^{-1}$ for all edges e ; and

- (iv) $E_{\mathcal{L}}$ is homotopy equivalent to the space $M_{\mathcal{L}}/f(\Gamma_{\mathcal{L}})$.

Section 3 contains proofs of Theorem 3 and Theorem 4. In Section 4 we show why the construction fails for general complex line arrangements and algebraic plane curves. The author thanks P. Orlik and M. Falk for helpful discussions during the writing of this paper.

2 Boundary manifolds

Let $\mathcal{L}$ be a line arrangement in the complex plane $\mathbb{C}^2$, let $\mathcal{A}$ be the set of lines in $\mathcal{L}$ and let $\mathcal{P}$ be the points of intersection on $\mathcal{L}$. In this section we describe the boundary manifold of $\mathcal{L}$ in terms of the incidence graph $\Gamma_{\mathcal{L}}$.

Case of disconnected incidence graph. Consider the case when $\mathcal{L}$ consists of k non-intersecting lines. This is the only case where $\Gamma_{\mathcal{L}}$ is not a connected graph and is a finite union of vertices, one for each line in $\mathcal{A}$. The boundary 3-manifold $M_{\mathcal{L}}$ is then a disjoint union of k solid tori. The complement $E_{\mathcal{L}}$ is equal to the product

$$\mathbb{C} \setminus \{k \text{ points}\} \times \mathbb{C}.$$

Thus, $M_{\mathcal{L}}/f(\Gamma_{\mathcal{L}})$ is homotopy equivalent to $E_{\mathcal{L}}$ for any map $f : \Gamma_{\mathcal{L}} \rightarrow M_{\mathcal{L}}$ which sends each v_L to a point on the line L .

Throughout this paper, unless otherwise stated we will assume that $\mathcal{P}$ is non-empty and therefore $\Gamma_{\mathcal{L}}$ is connected.

Incidence graph. The *(point/line) incidence graph* $\Gamma_{\mathcal{L}}$ of $\mathcal{L}$ is a bipartite graph with *point-vertices*

$$v_p, \quad p \in \mathcal{P}$$

and *line-vertices*

$$v_L, \quad L \in \mathcal{A}.$$

The edges of $\Gamma_{\mathcal{L}}$ are of the form

$$e(p, L) \text{ or } e(L, p), \quad p \in \mathcal{P}, L \in \mathcal{A}, \text{ and } p \in L.$$

The graph $\Gamma_{\mathcal{L}}$ is a directed graph. The *initial point* of $e = e(p, L)$ is defined to be $i(e) = v_p$ and the *terminal point* is defined to be $t(e) = v_L$. Similarly, if $e = e(L, p)$, then $i(e) = v_L$ and $t(e) = v_p$. We say that $e(L, p)$ and $e(p, L)$ are *conjugates* of each other and write $e(L, p) = \overline{e(p, L)}$.

We will also think of $\Gamma_{\mathcal{L}}$ as a singular 1-complex whose zero cells map to the vertices and whose one cells map to the edges so that e and $\bar{e}$ are identified but with opposite orientations. The endpoints of each edge e are attached to $i(e)$ and $t(e)$ in the obvious way.

Graph manifold. Let $\Gamma = \{\mathcal{V}, \mathcal{E}\}$ be a directed graph such that for each edge $e \in \mathcal{E}$, there is a conjugate edge $\bar{e} \in \mathcal{E}$ so that $i(e) = t(\bar{e})$ and $t(e) = i(\bar{e})$. A *graph manifold* $\mathcal{M}$ over Γ is a collection $(\{M_v\}, \{M_e\}, \{\phi_e\})$ where

- (i) M_v is a connected manifold with boundary for all $v \in \mathcal{V}$;
- (ii) M_e is a connected compact manifold without boundary for all $e \in \mathcal{E}$;
- (iii) $M_e = M_{\bar{e}}$ for all $e \in \mathcal{E}$;

(iv)

$$\phi_e : M_e \rightarrow M_{t(e)}$$

is an embedding of M_e onto a boundary component of $M_{t(e)}$;

(v) the map ϕ_e induces an endomorphism

$$\phi_{e*} : \pi_1(M_e) \rightarrow \pi_1(M_{t(e)})$$

on fundamental groups.

The underlying space M associated to a graph manifold $\mathcal{M}$ is defined to be the space obtained by gluing together the vertex manifolds M_v along their boundary components so that $\phi_e(M_e)$ is identified with $\phi_{\bar{e}}(M_e)$ by $\phi_e(q) = \phi_{\bar{e}}(q)$ for all $q \in M_e$. Any connect sum of manifolds glued along incompressible boundary components can be thought of as a graph of manifolds.

Graphs of groups. As with graph complexes, one can talk about the graph of groups associated to a graph manifold. A graph of groups $\mathcal{G}$ over a directed graph Γ is a collection of groups G_v , for each vertex $v \in \mathcal{V}$, groups $G_e = G_{\bar{e}}$, for each edge $e \in \mathcal{E}$, and group endomorphisms

$$\psi_e : G_e \rightarrow G_{t(e)}.$$

The collection of fundamental groups of the vertex and edge manifolds of a graph of manifolds is a graph of groups. The underlying group of a graph of groups is obtained from the vertex and edges groups by a combination of amalgamated products and HNN extensions. If $\mathcal{G}$ is the graph of groups associated to a graph of manifolds $\mathcal{M}$ then the underlying group of $\mathcal{G}$ is the fundamental group of the underlying manifold M of $\mathcal{M}$.

Regular neighborhood. Consider $\mathbb{C}^2$ as a metric space with the usual distance function

$$d((x_1, y_1), (x_2, y_2)) = \sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2}.$$

Let $\epsilon > 0$ be such that $d(p, L) > 2\epsilon$ for all pairs $(p, L) \in \mathcal{P} \times \mathcal{A}$ such that $p \notin L$.

For each $p \in \mathcal{P}$, let N_p be the ball of radius ϵ around p in $\mathbb{C}^2$ and let S_p be the boundary of N_p . Then N_p is a Milnor ball around p and the pair $(N_p, N_p \cap \mathcal{L})$ is a cone over $(S_p, S_p \cap \mathcal{L})$ (see [Mil]). Note that $L \cap N_p \neq \emptyset$ if and only if $p \in L$. Let

$$\delta_p = \min\{d(L_1 \cap S_p, L_2 \cap S_p)\}$$

where the minimum is taken over $L_1, L_2 \in \mathcal{A}$ and $p \in L_1 \cap L_2$. Take $\delta > 0$ such that $\delta_p > 2\delta$ for all $p \in \mathcal{P}$. For each line $L \in \mathcal{A}$, let

$$N_L = \{q \in \mathbb{C}^2 : d(q, L) \leq \delta\}.$$

By choosing δ smaller if necessary, we can assume that for each edge $e(p, L)$ in $\Gamma_{\mathcal{L}}$, N_L meets S_p transversally. Let

$$N_{\mathcal{L}} = \bigcup_{p \in \mathcal{P}} N_p \cup \bigcup_{L \in \mathcal{A}} N_L.$$

We call $N_{\mathcal{L}}$ a *regular neighborhood* of $\mathcal{L}$ in $\mathbb{C}^2$.

Boundary manifold. Consider the projective compactification $\mathbb{P}^2$ of $\mathbb{C}^2$ and let L_{∞} be the line at infinity, so that $\mathbb{P}^2 = \mathbb{C}^2 \cup L_{\infty}$. Let Q be a point on L_{∞} not on the projective closure $\overline{\mathcal{L}}$ of $\mathcal{L}$. Let $\overline{L}$ be the projective closure of L for all $L \in \mathcal{L}$.

For each $L \in \mathcal{A}$, let B_L be the closure of N_L in $\mathbb{P}^2$. Then B_L is a closed tubular neighborhood of $\overline{L}$ in $\mathbb{P}^2$. Let B_{∞} be a closed tubular neighborhood of L_{∞} not containing any element of $\mathcal{P}$. Let S_L, S_{∞} be the boundaries of B_L and B_{∞} , respectively.

The boundary $M_{\mathcal{L}}$ of $N_{\mathcal{L}}$ has a deformation retraction onto $M_{\mathcal{L}} \setminus B_{\infty}$. Hereafter, for simplicity, we will replace $M_{\mathcal{L}}$ by $M_{\mathcal{L}} \setminus B_{\infty}$.

For each $p \in \mathcal{P}$, note that

$$\bigcup_{L \in \mathcal{A}} N_L$$

intersects S_p transversally in disjoint 2-tori. Set

$$M_p = S_p \setminus \bigcup_{L \in \mathcal{A}} N_L.$$

For each $L \in \mathcal{A}$, let

$$M_L = S_L \setminus (B_{\infty} \cup \bigcup_{p \in \mathcal{P}} N_p).$$

Then $M_{\mathcal{L}}$ is a connect sum of the M_p and M_L .

By our construction we have the following.

Lemma 2.1 *The submanifolds M_p and M_L in $M_{\mathcal{L}}$ intersect in a boundary component if and only if $e(p, L)$ is an edge in $\Gamma_{\mathcal{L}}$, and this intersection is a 2-dimensional torus.*

Recall that a Hopf fibration $h : S^3 \rightarrow S^2$ is an oriented circle bundle such that the fibers have intersection number 1. Given any d points $\mathcal{Q}$ in S^2 , the preimage $h^{-1}(\mathcal{Q})$ is called a d -component Hopf link (see Figure 1). The pair $(S^3, h^{-1}(\mathcal{Q}))$ does not depend on which d points are chosen. Let H_d be a thickening of $h^{-1}(\mathcal{Q})$.




Figure 1. 3-component Hopf link.

Lemma 2.2 *For each $p \in \mathcal{P}$ there is a natural identification of the pair (S_p, M_p) with $(S^3, S^3 \setminus H_d)$, where d is the degree of v_p . For each $L \in \mathcal{A}$, there is a natural identification of the pair (S_L, M_L) with the pair $(S^3, S^3 \setminus H_{d+1})$, where d is the degree of v_L .*

Proof. We begin with M_p . Identify S_p with the 3-sphere S^3 and L_∞ with S^2 . Then the pencil of lines through p determines a projection $\rho : S_p \rightarrow L_\infty$ giving a commutative diagram

$$\begin{array}{ccc} S_p & \xrightarrow{=} & S^3 \\ \downarrow \rho & & \downarrow h \\ L_\infty & \xrightarrow{=} & S^2 \end{array}$$

where h is the standard Hopf fibration.




Figure 2. Neighborhood of a point $p \in \mathcal{P}$.

Let $L_1, \dots, L_d \subset \mathcal{L}$ be the lines passing through p . Under the identification of S_p with S^3 , M_p is identified with the complement in S^3 of

$$h^{-1}\left(\bigcup_{i=1}^d D_i\right),$$

where each D_i is a small disk around the point in S^2 corresponding to $\overline{L_i} \cap L_\infty$. We thus have a natural identification of the pair (S_p, M_p) with $(S^3, S^3 \setminus H_d)$.

Describing M_L is for the most part the dual picture to the above. Let S_L be the boundary of a tubular neighborhood of $S_{\overline{L}}$. Let B_Q be a ball neighborhood of Q in $\mathbb{P}^2$ and let S_Q be its boundary 3-sphere. Then S_L and S_Q are canonically identified as 3-spheres fibering over a 2-sphere. The pencil of lines through Q defines a commutative diagram

$$\begin{array}{ccc} S_Q & \xrightarrow{=} & S^3 \\ \downarrow \rho & & \downarrow h \\ \overline{L} & \xrightarrow{=} & S^2 \end{array}$$

where again h is the standard Hopf fibration. Under the identification of S_Q with S_L , B_∞ equals $h^{-1}(D_\infty)$ where D_∞ is a small disk neighborhood of $\overline{L} \cap L_\infty$.

Let $p_1, \dots, p_d \in \mathcal{P}$ be the points in $\mathcal{P}$ which lie on L . Then the identification of S_L with




Figure 3. Neighborhood of a line $L \in \mathcal{A}$.

S_Q gives an identification of M_L with

$$S_Q \setminus h^{-1}(D_\infty \cup \bigcup_{i=1}^d D_i),$$

where each D_i is a small disk neighborhood of p_i in $\overline{L}$. This identifies (S_L, M_L) with $(S^3, S^3 \setminus H_{d+1})$. ■

Lemma 2.1 and Lemma 2.2 imply the following.

Proposition 2.3 *For any line arrangement $\mathcal{L}$, $M_{\mathcal{L}}$ is a Haken 3-manifold with a torus decomposition into Seifert fibered manifolds. This torus decomposition gives $M_{\mathcal{L}}$ the structure of a graph manifold over $\Gamma_{\mathcal{L}}$.*

Proof. We only need to check when the torus boundary components of $S^3 \setminus H_d$ are incompressible. This is true as long as $d > 1$, which holds unless $\mathcal{L}$ is a union of parallel lines. ■

Corollary 2.4 *The fundamental group $\pi_1(M_{\mathcal{L}})$ is torsion free and residually finite.*

Proof. The fundamental group of $\pi_1(M_{\mathcal{L}})$ is a graph of groups whose vertex groups are $\pi_1(S^3 \setminus H_d) = \mathbb{Z} \times F_d$, where F_d is a free group on $d - 1$ generators, and are torsion free. It follows that $\pi_1(M_{\mathcal{L}})$ is also torsion free (see, for example, Section 1.3, Corollary 2 and Section 1.4, Proposition 5 of [Ser].)

Since $M_{\mathcal{L}}$ is a 3-manifold with a torus decomposition along incompressible tori into Seifert fibered pieces, by [Hemp], Theorem 1.1, the fundamental is residually finite. ■

The gluing maps.

Let T be a 2-dimensional torus. A *framing* (μ, λ) on T is a choice of generators for its fundamental group $\pi_1(T)$.

As stated in Proposition 2.3, the edge manifolds of $M_{\mathcal{L}}$ considered as a graph manifold are all 2-dimensional tori, so they are isomorphic to T . To prove Theorem 2, we need to describe the gluing maps

$$\phi_e : T \rightarrow M_p$$

and

$$\phi_{\bar{e}} : T \rightarrow M_L$$

for each edge $e = e(p, L)$ on the graph $\Gamma_{\mathcal{L}}$.

By Lemma 2.2, the vertex manifolds M_p and M_L are naturally identified with $E_d = S^3 \setminus H_d$ for some integer $d \geq 2$. Furthermore, the complex structure on M_p and M_L determines an orientation on the core loops of H_d . The oriented pair (S^3, H_d) determines a framing on the boundary components of E_d and hence on the boundary components of M_p and M_L .

Let T be a boundary component of E_d and let ℓ be its core curve in S^3 . The framing (μ, λ) on T is given as follows. Choose a basepoint x on T . The loop μ is a positively oriented meridian loop around the core curve based at x . If one considers T as an S^1 -bundle over ℓ , then μ is a loop going once in the positive direction around the fiber of the bundle. The loop λ is a loop based at x “parallel” to the core curve. That is, it is a positively oriented loop whose linking number with the core curve is zero. These definitions uniquely determine μ and λ up to homotopy.

Let $p \in \mathcal{P}$, $L \in \mathcal{A}$ be such that $p \in L$. We will write $T_{L,p}$ for the boundary component on M_p corresponding to L , and $T_{p,L}$ for the boundary component on M_L corresponding to p . Let $(\mu_{L,p}, \lambda_{L,p})$ be the framing on $T_{L,p}$ and $(\mu_{p,L}, \lambda_{p,L})$ be the framing on $T_{p,L}$.

To describe ϕ_e up to homotopy, it suffices to describe what it does to the framings.

Lemma 2.5 *Let $e = e(L, p)$ be an edge of $\Gamma_{\mathcal{L}}$. If we identify T with $T_{L,p}$ by the map*

$$\begin{aligned} \phi_e : T &\rightarrow T_{L,p} \subset M_p \\ \phi_e(\mu) &= \mu_{L,p} \\ \phi_e(\lambda) &= \lambda_{L,p}, \end{aligned}$$

then

$$\begin{aligned} \phi_{\bar{e}} : T &\rightarrow T_{p,L} \subset M_L \\ \phi_{\bar{e}}(\mu) &= \mu_{p,L} + \lambda_{p,L} \\ \phi_{\bar{e}}(\lambda) &= \mu_{p,L} \end{aligned}$$

gives the gluing map for the graph manifold $M_{\mathcal{L}}$.




Figure 4. Intersection of S_L and S_p .

Proof. We will study how S_L and S_p meet by changing coordinates so that L is given by the equation $y = 0$, the point p is the origin, $p = (0, 0)$. Then S_L and S_p are given by

$$S_L = \overline{\{(x, y) \in \mathbb{C}^2 : |y| = 1\}},$$

and

$$\begin{aligned} S_p &= \{(x, y) \in \mathbb{C}^2 : |x| \leq 1, |y| = 1\} \cup \{(x, y) \in \mathbb{C}^2 : |x| = 1, |y| \leq 1\} \\ &= \mathbb{T}_1 \cup \mathbb{T}_2 \end{aligned}$$

as shown in Figure 4. Then $S_L \cap S_p = T$, where T is the 2-torus

$$T = \{(x, y) \in \mathbb{C}^2 : |x| = 1, |y| = 1\}$$

and is the common boundary of $\mathbb{T}_1$ and $\mathbb{T}_2$.

As before let $T_{L,p}$ be the boundary component of S_p corresponding to L and $T_{p,L}$ be the boundary component of S_L corresponding to p . Since the components of the Hopf link are all unknots, the framing on the link boundary components is given as follows. Each link component has a solid torus neighborhood N in S^3 . The complement of N is also a solid torus N^c . The framing on the boundary of N is given by (μ, λ) where μ contracts in N and λ contracts in N^c .

Thus, $T_{L,p}$ has the framing

$$\begin{aligned} \mu_{L,p} &= (1, \exp(2\pi it)) \\ \lambda_{L,p} &= (\exp(2\pi it), 1) \end{aligned}$$

and $T_{p,L}$ has the framing

$$\begin{aligned} \mu_{p,L} &= (\exp(2\pi it), 1) \\ \lambda_{p,L} &= (\exp(2\pi it), \exp(2\pi it)). \end{aligned}$$

This implies $\mu_{p,L} = \lambda_{L,p}$ and $\lambda_{p,L} = \mu_{L,p} + \lambda_{L,p}$, as illustrated in Figure 5.

■




Figure 5. Gluing map.

Proof of Theorem 2.

We have shown that $M_{\mathcal{L}}$ is a graph manifold whose vertex manifolds are $S^3 \setminus H_d$, where d is the degree of the vertex if the vertex is a point-vertex and $d - 1$ is the degree of the vertex if the vertex is a line-vertex. The identifications of the edge manifolds with the boundary components of the vertex manifolds can be described in terms of the natural framings on the boundary components of $S^3 \setminus H_d$ by the edge maps described in Lemma 2.5. This completes the proof of Theorem 2. ■

3 Real line arrangements

In this section, we concentrate on real line arrangements, and will assume that $\mathcal{L}$ is defined by equations of the form

$$y = a_i x + b_i, \quad i = 1, \dots, k,$$

where $a_i, b_i \in \mathbb{R}$. We will show how to reconstruct the homotopy type of the complement of $\mathcal{L}$ in terms of the ordered graph of $\mathcal{L}$.

Ordered graphs.

The ordered graph associated to $\mathcal{L}$ is the incidence graph $\Gamma_{\mathcal{L}}$ together with some extra structure. For real line arrangements, we will order the edges emanating from each vertex of the incidence graph $\Gamma_{\mathcal{L}}$ as follows. If v_p is the point-vertex associated to the point $p \in \mathcal{P}$, then we order the edges $e(p, L_1), \dots, e(p, L_r)$ emanating from p by the slopes of $L_1, \dots, L_r$ in decreasing order. If v_L is the line-vertex associated to the line $L \in \mathcal{A}$, we order the edges $e(L, p_1), \dots, e(L, p_s)$ emanating from v_L by the x -coordinates of $p_1, \dots, p_s$ in decreasing order. Since we assume that none of the lines are parallel to the y -axis, this is well-defined. The incidence graph $\Gamma_{\mathcal{L}}$ of $\mathcal{L}$ endowed with these orderings on the edges emanating from vertices is called the *ordered graph* associated to $\mathcal{L}$. For example, Figure 6 gives the ordered graph of the Ceva arrangement.




Figure 6. Ordered graph associated to the Ceva arrangement.

The global ordering of the points in $\mathcal{P}$ by their x -coordinate and the lines in $\mathcal{A}$ by their slope determines the ordered graph. The orderings near v_{L_1} and v_{p_5} are given in Figure 6.

The skeleton of a line arrangement.

Let $M_{\mathcal{L}}$ be the boundary manifold of $\mathcal{L}$ and let

$$\alpha : M_{\mathcal{L}} \rightarrow \mathcal{L}$$

be the natural projection map, well-defined up to homotopy. Let $\Sigma_{\mathcal{L}}$ be the real part of $\mathcal{L}$ with all infinite ends removed. We will call $\Sigma_{\mathcal{L}}$ the skeleton of $\mathcal{L}$. Figure 7 gives the skeleton of the Ceva arrangement.




Figure 7. Skeleton associated to the Ceva arrangement.

We start by describing the homotopy type of $E_{\mathcal{L}}$ in terms of $M_{\mathcal{L}}$ and $\Sigma_{\mathcal{L}}$.

Lemma 3.1 *There is an embedding*

$$g : \Sigma_{\mathcal{L}} \rightarrow M_{\mathcal{L}}$$

so that

- (i) $\alpha \circ g$ is the identity map;
- (ii) the image of g contracts in $E_{\mathcal{L}}$; and
- (iii) $E_{\mathcal{L}} = M_{\mathcal{L}}/g(\Sigma_{\mathcal{L}})$.

Proof. The embedding g is given as follows. If $(x, y) \in N_p$, for some $p = (x_p, y_p) \in \mathcal{P}$, then

$$g(x, y) = (x, y + i \max\{\sqrt{\epsilon^2 - (y - y_p)^2}, \delta\}),$$

otherwise

$$g(x, y) = (x, y + i\delta).$$

The image $g(\Sigma_{\mathcal{L}})$ contracts in $E_{\mathcal{L}}$, since in $E_{\mathcal{L}}$ it is isotopic to $\Sigma_{\mathcal{L}} + (0, i\epsilon) \subset \mathbb{R}^2 + (0, i\epsilon)$. No point $(x, y + i\epsilon) \in \mathbb{R}^2 + (0, i\epsilon)$ satisfies an equation of the form

$$y = ax + b$$

where a and b are real numbers. Thus, $\mathbb{R}^2 + (0, i\epsilon)$ is contained in $E_{\mathcal{L}}$. Since $\mathbb{R}^2 + (0, i\epsilon)$ is contractible, we see that $g(\Sigma_{\mathcal{L}})$ is also contractible.

Now let us consider the projection of $\rho : \mathbb{C}^2 \rightarrow \mathbb{C}$ onto the first coordinate. Let $\rho_{\mathcal{L}}$ be the restriction of ρ to $E_{\mathcal{L}}$. Then $\rho_{\mathcal{L}}$ is a topological fibration over the complement of $\mathcal{Q} = \rho(\mathcal{P})$.




Figure 8. Contraction of the x -coordinate plane to V_Q .

For each $q \in \mathcal{Q}$ let

$$D_q = \{x \in \mathbb{C} : |x - q| \leq 1\}$$

and let S_q be the boundary of D_q . Assume (by expanding coordinates if necessary) that the S_q do not intersect one another. Let W_Q be the union of the S_q and real line segments joining the S_q , and let V_Q be the union of W_Q and the D_q as in Figure 8.

The deformation retraction of $\mathbb{C}$ onto V_Q extends to a deformation retraction of $E_{\mathcal{L}}$ onto $\rho_{\mathcal{L}}^{-1}(V_Q)$.

The fibers of $\rho_{\mathcal{L}}$ over V_Q split up into 3 types: those over the line segments in W_Q ; those over S_q ; and those over the q . These retract to the outlined and shaded regions shown in Figure 9 which we will write as F_x , for $x \in I$, F_s , for $s \in S$ and F_q , respectively.

We reconstruct the homotopy type of $E_{\mathcal{L}}$ from this picture as follows. First notice that the space $\rho^{-1}|_{M_{\mathcal{L}}}(D_q)$ has a deformation retraction to

$$F_{D_q} = D_q \times F_q \cup \bigcup_{s \in S_q} F_s.$$




Figure 9. Fibers over V_Q .

Under this retraction $g(\Sigma_{\mathcal{L}})$ maps to the right most points of the circles in F_s , F_x and F_q union the line segments depicted on F_{s_1} and F_{s_2} . The union of the right-most points of the circles on F_s as s ranges in S_q bounds a 2-disk

$$\{p + (0, iy) : |y| \leq \epsilon\} = D_q \times \{p + (0, i\epsilon)\}$$

in the deformation of $\rho^{-1}|_{\mathcal{L}}(D_q)$. Thus, if we include all right-most points of circles in the fibers over F_x , F_s , and F_q , union the line segments on F_{s_1} and F_{s_2} , we obtain a set G which retracts onto $g(\Sigma_{\mathcal{L}})$ in $E_{\mathcal{L}}$. ■

Proof of Theorem 4.

The existence of a map $f : \Gamma_{\mathcal{L}} \rightarrow M_{\mathcal{L}}$ with property (iii) of Theorem 4 follows from Lemma 3.1, since $\Gamma_{\mathcal{L}}$ can be continuously deformed to $\Sigma_{\mathcal{L}}$. One can see this by noting that the figures in Figure 10 are homotopy equivalent.

We will explicitly describe the homotopy type of a map $f : \Gamma_{\mathcal{L}} \rightarrow M_{\mathcal{L}}$ satisfying the conditions of Theorem 4 in terms of the ordered graph.

We begin by defining a map $\sigma : \Gamma_{\mathcal{L}} \rightarrow \Sigma_{\mathcal{L}}$. For the point-vertices, we simply send v_p to the point p on $\Sigma_{\mathcal{L}}$. For the line vertices, there is no canonical choice.

Let $L \in \mathcal{A}$. If $\text{degree}(v_L) = 1$, then let $\sigma(v_L) = p$ where $p = \mathcal{P} \cap L$. Otherwise, let $p_1, p_2, \dots, p_r$ be the points in $\mathcal{P} \cap L$. Choose a point on the line segment on $\Sigma_{\mathcal{L}}$ strictly between p_1 and p_2 and map v_L to this point.




Figure 10. Incidence graph and skeleton.

For the edges of $\Gamma_{\mathcal{L}}$, once we have determined where the vertices go, the edges map to the unique straight line segment on $\Sigma_{\mathcal{L}}$ connecting the endpoints. This defines a continuous map $\sigma : \Gamma_{\mathcal{L}} \rightarrow \Sigma_{\mathcal{L}}$. The composition $f = g \circ \sigma$ gives a map satisfying properties (i),(iii) and (iv) of Theorem 4.

For each vertex v on Γ , let $z_v = f(v)$. (We will write $z_p = z_{v_p}$ and $z_L = z_{v_L}$.) To describe the homotopy type of the map f , it suffices to describe the homotopy type of $f(e)$ for each edge e in Γ . The path $\sigma(e)$ breaks up into segments each passing over one point $p \in \mathcal{P}$. Let $I_{L,p}$ be a straight line segment on $\Sigma_{\mathcal{L}} \cap L$ such that $I_{L,p} \cap \mathcal{P} = p$. We will describe the homotopy type of $g(I_{L,p})$.




Figure 11. Path γ_L lifting to L_0 .

First fix L . Consider the path γ_L on the complex plane shown in Figure 11. Here $\mathcal{Q} = \{q_1, \dots, q_s\} = \rho(\mathcal{P} \cap L)$. The projection map ρ is one-to-one when restricted to L . Since $\rho^{-1}(\gamma_L) \cap L$ is contained in L_0 , $b_L = \rho^{-1}(\gamma_L) \cap L$ defines a contractible subset of L_0 . Let $h : \gamma_L \rightarrow M_L$ be defined by

$$h(x, y) = \rho|_L^{-1}(x, y) + (0, \delta i),$$

for each $(x, y) \in \gamma_L$. Then $h(\gamma_L)$ defines a contractible subset of M_L . We can assume that the image contains the basepoint z_L .




Figure 12. Generators for M_p .

Now fix $p \in L \cap \mathcal{P}$. Let $q = \rho(p)$ and let J_p be the arc segment of γ_L near q . For each fiber F_s over points $s \in J_p$, $h(J_p) \cap F$ is the rightmost point in the circle (see Figure 13) on F corresponding to $L \cap F$. As noted in the proof of Lemma 3.1, the path $g(I_p)$ is isotopic to the path whose intersection with each fiber F_s over an interior point s of J_p is the right-most point of the large circle and whose intersection with the fibers F_{s_1} and F_{s_2} is a path from the right-most point of the large circle to the right-most point of the inner circle associated to L (see Figure fig:fibers). The large disk in the fiber F_s over $s \in J_p$ rotates in the counter-clockwise direction by 180 degrees as s moves from right to left on J_p , as illustrated in Figure 13.




Figure 13. Homotopy type of $f(e(p, L))$.

Note that the basepoint z_p for M_p is the right-most point on the large circle on F_{s_1} and that $b_L \cap F_s$ is the right-most point on the inner circle corresponding to L for all $s \in J_p$. Let $\tau_{L,p}$ be the straight line segment from $b_L \cap F_{s_1}$ to z_p (see Figure 13.)

Let $\mu_{L_1,p}, \dots, \mu_{L_r,p}$ be the generators for $\pi_1(M_p)$ as in Figure 12. Then $g(I_p)h(J_p)^{-1}$ is homotopy equivalent to the path abc on F_{s_1} pictured in Figure 14. In this example L_1, L_2, L_3 are the lines passing through p in the order of their slopes and $L = L_2$, then $h(J_p)$ is homotopy equivalent to $g(I_p)h(J_p)^{-1} = abc = \tau_{L,p}\mu_{L_3,p}^{-1}\mu_{L_2,p}\mu_{L_3,p}\tau_{L,p}^{-1}$.

In the general case, we have the following lemma.

Lemma 3.2 *Let $p \in \mathcal{P}$ and let I_p be a line segment on $\Sigma_{\mathcal{L}} \cap L$ such that $I_p \cap \mathcal{P} = \{p\}$ and p lies in the interior of I_p . Let $L_1, \dots, L_r$ be an ordering by slope of the lines in $\mathcal{A}$ passing through p . Let j be such that $L_j = L$. Then the homotopy class of the difference $g(I_p)h(J_p)^{-1}$ on $M_{\mathcal{L}}$ is given by the element*

$$g_{L,p} = \tau_{L,p}\mu_{L_r,p}^{-1}\mu_{L_{r-1},p}^{-1} \cdots \mu_{L_j,p}\mu_{L_{j+1},p} \cdots \mu_{L_r,p}\tau_{L,p}^{-1}.$$




Figure 14. Homotopy type of $g(I_p)h(J_p)^{-1}$

We now describe the homotopy type of the map

$$f : \Gamma_{\mathcal{L}} \rightarrow M_{\mathcal{L}}$$

in terms of the ordered graph $\Gamma_{\mathcal{L}}$.

Fix $p \in \mathcal{P}$ and let $e(p, L_1), \dots, e(p, L_r)$ be the ordered edges emanating from v_p . For each line $L \in \mathcal{A}$, let $e(L, p_1), \dots, e(L, p_s)$ be the ordered edges emanating from v_L .

For any edge $e(L, p)$ in $\Gamma_{\mathcal{L}}$, replace $\tau_{L,p}$ by the path from $z_L \in M_L$ to $z_p \in M_p$ which goes along $h(\gamma_L)$ and the original $\tau_{L,p}$. This does not change the homotopy type of $\tau_{L,p}$. Note that the choices of $\tau_{L,p}$, z_p , z_L a continuous map

$$\ell : \Gamma_{\mathcal{L}} \rightarrow M_{\mathcal{L}}$$

where $\ell(v) = z_v$, for vertices v and $\ell(e(L, p)) = \tau_{L,p}$ for edges $e(L, p)$. We will call this map ℓ the *lifting* of $\Gamma_{\mathcal{L}}$ in $M_{\mathcal{L}}$ defined by the collection $\tau_{L,p}$. Let $g_{L,p} \in \pi_1(M_p, z_p)$ be defined as in Lemma 3.2, except that its basepoint is at z_L . We have shown the following.

Lemma 3.3 *Let*

$$f : \Gamma_{\mathcal{L}} \rightarrow M_{\mathcal{L}}$$

be the continuous map on $\Gamma_{\mathcal{L}}$ considered as a 1-complex defined by the following data:

- (i) *for the point-vertex v_p , $f(v_p) = z_p$;*
- (ii) *for the line-vertex v_L , $f(v_L) = z_L$; and*
- (iii) *for the edge $e(L, p)$, if $e(L, p_1), \dots, e(L, p_s)$ are the ordered edges emanating from v_L , and $p = p_j$, then*

$$f(e(L, p)) = \begin{cases} \tau_{L,p} & \text{if } j = 1, 2; \\ g_{L,p_2} \cdots g_{L,p_{j-1}} \tau_{L,p} & \text{if } j > 2 \end{cases}$$

and $f(e(p, L)) = f(e(L, p))^{-1}$. Then $E_{\mathcal{L}}$ is homotopy equivalent to $M_{\mathcal{L}}/f(\Gamma_{\mathcal{L}})$.

For each pair $(p_0, L_0) \in \mathcal{P} \times \mathcal{A}$ with $p_0 \in L_0$, the element g_{L_0,p_0} depends on the ordering of the edges $e(p_0, L)$ emanating from v_{p_0} and $f(e(L_0, p_0))$ depends on the ordering the edges $e(L_0, p)$ emanating from v_{L_0} . This completes the proof of Theorem 4. ■

Fundamental group.

Choose a vertex v of $\Gamma_{\mathcal{L}}$. We present the fundamental group $G_{\mathcal{L}} = \pi_1(M_{\mathcal{L}}, f(v))$ in the notation of [Ser] (p. 41). For each edge $e = e(L, p)$ in $\Gamma_{\mathcal{L}}$, let $\tau_e = \tau_{L,p}$ and let $\tau_{\bar{e}} = \tau_e^{-1}$. The corresponding lifting of $\Gamma_{\mathcal{L}}$ into $M_{\mathcal{L}}$ determines a presentation of $\pi_1(M_{\mathcal{L}})$ as follows.

Any element of $G_{\mathcal{L}}$ can be written as a word

$$\gamma_1 \tau_{e_1} \cdots \gamma_k \tau_{e_k} \gamma_{k+1} \quad (1)$$

satisfying

- (i) $t(e_i) = i(e_{i+1})$, for $i = 1, \dots, k$;
- (ii) $t(e_k) = i(e_1) = v$;
- (iii) each γ_i is an element of $\pi_1(M_{t(e_i)})$.

The relations on $G_{\mathcal{L}}$ are generated by elements of the form

$$\tau_e \gamma \tau_{\bar{e}} (\gamma')^{-1},$$

where $\gamma = \phi_e(\kappa)$ and $\gamma' = \phi_{\bar{e}}(\kappa)$ for some $\kappa \in G_e$. A word of the form (1) is said to have length k . The length thus depends on the word and not its equivalence class in $G_{\mathcal{L}}$. A word of the form (1) is *reduced* if whenever $e_i = \overline{e_{i+1}}$, for some $i = 1, \dots, k-1$, then $\gamma_{i+1} \notin \phi_{e_i}(\pi_1(M_{e_i}))$.

The following theorem is useful for determining properties of groups which are graphs of groups.

Proposition 3.4 ([Ser], Theorem 11.) *A reduced word of the form (1) is trivial if and only if it has length 1 and $\gamma_1 = 1$.*

For example, we have the following result.

Corollary 3.5 *The map $f : \Gamma_{\mathcal{L}} \rightarrow M_{\mathcal{L}}$ induces an endomorphism on fundamental groups.*

Proof. Let $G_{\mathcal{L}}$ be the image of

$$f_* : \pi_1(\Gamma_{\mathcal{L}}, v) \rightarrow \pi_1(M_{\mathcal{L}}, f(v)).$$

Then $G_{\mathcal{L}}$ consists of elements of the form

$$f(e_1) \cdots f(e_k) = f_*(e_1 \cdots e_k) \quad (2)$$

where $e_1 \cdots e_k$ is a closed loop on $\Gamma_{\mathcal{L}}$ and $i(e_1) = t(e_k) = v$. Suppose an element of the form (2) is trivial in $G_{\mathcal{L}}$ and $k > 1$. We will show that we can write the same element in the form (2) with smaller k . By Theorem 5, $k \geq 2$ and there is at least one i so that $e_i = \overline{e_{i+1}}$. In this case $e_i e_{i+1} = 1$, so the element can be represented with smaller k . ■

We will now present the fundamental groups of $M_{\mathcal{L}}$ and of $E_{\mathcal{L}}$ purely from the ordered graph $\Gamma_{\mathcal{L}}$.

Take any ordered bipartite graph $\tilde{\Gamma}$ with “line” and “point” vertices, such that each directed edge $e(p, L)$ (or $e(L, p)$) joins a point-vertex v_p to a line-vertex v_L (or vice versa.) Choose a vertex v_0 on $\tilde{\Gamma}$. For v , a vertex on $\tilde{\Gamma}$, let $e_{v,1}, \dots, e_{v,d}$ be the ordering of edges emanating from v . Set

$$G_v = \langle \lambda_v, \mu_{1,v}, \dots, \mu_{d,v} : \lambda_v \mu_{j,v} = \mu_{j,v} \lambda_v, \quad j = 1, \dots, d \rangle.$$

(Note that in the previous notation $\lambda_{j,v} = \lambda_v \mu_{j,v}$, for each $j = 1, \dots, \text{degree}(v)$.) Let $\mathcal{T}_{\Gamma}$ be the collection of closed loops on Γ based at v_0 . Then, for any $\tau \in \mathcal{T}_{\Gamma}$, $\tau = e_1 \dots e_r$, where $t(e_j) = i(e_{j+1})$, for $j = 1, \dots, r-1$ and $i(e_1) = t(e_r) = v_0$.

The following is a corollary of Theorem 2.

Corollary 3.6 *Let $\mathcal{L}$ be a complex line arrangement. Then $\pi_1(M_{\mathcal{L}})$ can be presented as follows: each element of $\pi_1(M_{\mathcal{L}})$ can be written as*

$$\gamma_0 e_1 \gamma_1 e_2 \dots e_s \gamma_s,$$

where $e_1 \dots e_s \in \mathcal{T}_{\Gamma_{\mathcal{L}}}$ and $\gamma_j \in G_{t(e_j)}$, for $j = 1, \dots, s$, and the relations are generated by

$$\begin{aligned} \alpha e(p, L) \mu_{k,v_L} e(L, p) \beta &= \alpha \lambda_{v_p} \mu_{j,v_p}^{-1} \beta, \\ \alpha e(p, L) \lambda_{v_L} \mu_{k,v_L}^{-1} e(L, p) \beta &= \alpha \lambda_{v_p} \beta, \end{aligned}$$

if $e(p, L)$ is the j th edge emanating from v_p , and $e(L, p)$ is the k th edge emanating from v_L .

Note that the indexing of the edges at each vertex does not effect the isomorphism class of the group presented.

By Theorem 4, $\pi_1(E_{\mathcal{L}})$ is a quotient of $\pi_1(M_{\mathcal{L}})$ and we have the following presentation of $\pi_1(E_{\mathcal{L}})$.

Corollary 3.7 *Let $\mathcal{L}$ be a real line arrangement. Then the fundamental group of the complement $\pi_1(E_{\mathcal{L}})$ is $\pi_1(M_{\mathcal{L}})$ with the additional relations*

$$f(e_1) \dots f(e_s) = 1$$

where f is as in Theorem 4 and $e_1 \dots e_s \in \mathcal{T}_{\Gamma}$.

4 Complex line arrangements

Let $\mathcal{L}$ be an arbitrary complex line arrangement consisting of lines $\mathcal{A}$ and points of intersection $\mathcal{P}$. The proofs of Theorem 3 and Theorem 4 given in Section 3 rely on the existence of a continuous map

$$g : \Sigma_{\mathcal{L}} \rightarrow M_{\mathcal{L}}$$

from the skeleton $\Sigma_{\mathcal{L}}$ of $\mathcal{L}$ to the boundary 3-manifold $M_{\mathcal{L}}$ whose image contracts in $E_{\mathcal{L}}$.

We will show that Theorem 3 and Theorem 4 cannot be applied to arbitrary complex line arrangements. To do this we extend the definition of skeleton for complex line arrangements, and show that the map g and hence the map

$$f : \Gamma_{\mathcal{L}} \rightarrow M_{\mathcal{L}}$$

with the required properties does not exist in general.

Knotted 1-complexes.

Let $\Sigma \subset \mathbb{R}^3$ be a compact 1-complex. We will call Σ a *graph knot*. Let N_{Σ} be a thickening of Σ in $\mathbb{R}^3$ and let M_{Σ} be its boundary. Then N_{Σ} is a handlebody and M_{Σ} is an oriented surface. Let $\alpha' : N_{\Sigma} \rightarrow \Sigma$ be the contraction map and let $\alpha : M_{\Sigma} \rightarrow \Sigma$ be the restriction of α' to M_{Σ} . We say Σ is *unknotted* if there is an embedding

$$g : \Sigma \rightarrow M_{\Sigma}$$

such that $\alpha \circ g$ is the identity on Σ , and the image of g contracts in $\mathbb{R}^3$. We say that Σ is *knotted* otherwise.

Dehn's Lemma ([Rolf], p. 101) implies the following.

Lemma 4.1 *A graph knot $\Sigma \subset \mathbb{R}^3$ is knotted if there is an embedding*

$$\sqcup S^1 \rightarrow \Sigma$$

whose image is a nontrivial knot or link in $\mathbb{R}^3$.

Skeleta for complex arrangements.

Let $\mathcal{L}$ be an arbitrary complex line arrangement and let

$$\rho : \mathbb{C}^2 \rightarrow \mathbb{C}$$

be projection onto the x -coordinate. Assume that $\rho|_L$ is one-to-one for all lines $L \in \mathcal{A}$. Let $\mathcal{Q} = \rho(\mathcal{P})$.

Given an ordering of the points $q_1, \dots, q_s \in \mathcal{Q}$, let $I = I_1 \cup \dots \cup I_{s-1}$ be a union of embedded line segments in $\mathbb{C}$ such that each I_i has endpoints q_i and q_{i-1} and no pair I_i and I_j intersect except possibly at their endpoints. We will call $\Sigma_{\mathcal{L}}(I) = \rho^{-1}(I) \cap \mathcal{L}$ the *skeleton* of $\mathcal{L}$ associated to I . (The 1-complex $\Sigma_{\mathcal{L}}(I)$ considered as a subset of $\mathbb{R}^2 \times I$ is also known as a *braided wiring diagram* and has been used to describe the fundamental group and homotopy type of the complement of arbitrary complex line arrangements and algebraic plane curves [Arv], [C-S].)

The skeleton $\Sigma_{\mathcal{L}}$ defined for real line arrangements comes from ordering the q_i in the order in which they lie on the real line $\mathbb{R}$ and taking I to be the line segment on $\mathbb{R}$ connecting the smallest to the largest. We will call this $\Sigma_{\mathcal{L}}$ the *standard skeleton* for the real line arrangement $\mathcal{L}$.

As with real line arrangements, we have the following.

Lemma 4.2 *For any complex line arrangement $\mathcal{L}$ and any ordering of $\mathcal{Q}$, the 1-complex $\Sigma_{\mathcal{L}}(I)$ is homotopy equivalent to the graph $\Gamma_{\mathcal{L}}$.*

We can think of $\Sigma_{\mathcal{L}}(I)$ as a subset of $\mathbb{R}^3 = \mathbb{R}^2 \times \mathbb{R}$ by embedding $I \in \mathbb{R}$. This makes $\Sigma_{\mathcal{L}}(I)$ a graph knot. When $\mathcal{L}$ is a real line arrangement and $\Sigma_{\mathcal{L}}$ is the standard skeleton $\Sigma_{\mathcal{L}}$ is unknotted. Conversely, if $\mathcal{L}$ is a complex line arrangement such that for all orderings the associated skeleton $\Sigma_{\mathcal{L}}(I)$ is knotted, then $\mathcal{L}$ is not topologically equivalent to a real line arrangement.

Lemma 3.1 then generalizes as follows.

Theorem 5 *Let $\mathcal{L}$ be a complex line arrangement and let $\alpha : M_{\mathcal{L}} \rightarrow \mathcal{L}$ be the natural projection map of the boundary manifold $M_{\mathcal{L}}$ onto $\mathcal{L}$. Then for some choice of I , $\Sigma_{\mathcal{L}}(I)$ is unknotted if and only if there is an embedding*

$$g : \Sigma_{\mathcal{L}}(I) \rightarrow M_{\mathcal{L}},$$

such that

(i) $\alpha \circ g$ is the identity map; and

(ii) the complement $E_{\mathcal{L}}$ is homotopy equivalent to

$$M_{\mathcal{L}}/g(\Sigma_{\mathcal{L}}(I)).$$

Proof. The proof is the same as for Lemma 3.1. ■

It follows that if $\Sigma_{\mathcal{L}}(I)$ is knotted for all choices of I , then Theorem 4 cannot be extended to $\mathcal{L}$. For general algebraic plane curves, one can also describe the boundary 3-manifold $M_{\mathcal{C}}$ as a graph of manifolds over a suitably defined incidence graph $\Gamma_{\mathcal{C}}$, but Theorem 5 provides an obstruction for mimicing Theorem 4 in this setting.

References

- [Arv] W. Arvola. The fundamental group of the complement of an arrangement of complex hyperplanes. *Topology* **31**(1992), 757–765.
- [C-S] D. Cohen and A. Suciu. Braided monodromy of plane algebraic curves and hyperplane arrangements. *Comm. Math. Helv.* **72**(1997), 285–315.
- [Falk] M. Falk. Homotopy Types of Line Arrangements. *Invent. Math.* **111**(1993), 139–150.
- [G-M] M. Goresky and R. Macpherson. *Stratified Morse Theory*, volume 14 of *Ergebnisse der Mathematik und ihrer Grenzgebiete (3)*. Springer-Verlag, Berlin, 1988.

- [Hemp] J. Hempel. Residual finiteness for 3-manifolds. In *Combinatorial group theory and topology (Alta, Utah, 1984)*, volume 111 of *Ann. of Math. Stud.*, pages 379–396. Princeton University Press, 1987.
- [Lib] A. Libgober. On the homotopy type of the complement to plane algebraic curves. *J. Reine Angew. Math.* **367**(1986), 103–114.
- [Mil] J. Milnor. *Singular Points of Complex Hypersurfaces*, volume 61 of *Annals of Math. Studies*. Princeton Univ. Press, Princeton, 1968.
- [O-T] P. Orlik and H. Terao. *Arrangement of Hyperplanes*, volume 300 of *Grundlehren der math. Wissenschaften*. Springer-Verlag, Berlin, 1992.
- [Pen] R. Penner. Perturbative series and the moduli space of Riemann surfaces. *J. Diff. Geo.* **27**(1988), 35–53.
- [Ran] R. Randell. The fundamental group of the complement of a union of complex hyperplanes (Correction). *Invent. Math.* **69 (80)**(1982 (1985)), 103–108 (467–468).
- [Rolf] Rolfsen. *Knots and Links*.
- [Ryb] G. Rybnikov. On the fundamental group of the complement of a complex hyperplane arrangement. *DIMACS: Technical Report* (1994), 33–50.
- [Sal] M. Salvetti. Topology of the complement of real hyperplanes in $\mathbb{C}^N$. *Invent. Math.* **88**(1987), 603–618.
- [Ser] J-P. Serre. *Trees*. Springer-Verlag, Berlin, 1980.
- [Wal] F. Waldhausen. Eine Klasse von 3-dimensionalen Mannigfaltigkeiten I. *Invent. Math.* **3**(1967), 308–333.